



PROTOCOL 3000 with DMSO

The 3000 protocol is often confused with the **AMS** bottle protocol for external application of **AMS**. The 3000 protocol is used to allow the **AMS** along with **DMSO** to be absorbed through the skin penetrating tumors entering the blood and even penetrating bone.

The goal with life-threatening situations is to quickly get **AMS** circulating in the blood while trying to stay under the “nausea level”. One way to achieve this is by adding **DMSO** to activated **AMS**, so that it can act as a carrier for sending **AMS** directly into the skin and muscles and thus into the blood. Experience and testing have proven that **DMSO** is carried directly to any cancer in the body and it then penetrates the cancer cells. This is not speculation; it has been proven through testing. The theory is that when **DMSO** is carrying **AMS**, it will take it into the cancerous cells, thereby killing the virus that makes the cells cancerous. Many cancers have been healed in this way, and I hear about it since many people have phoned me stating that their cancer is gone.

- When using this treatment, always also take **AMS** by mouth, as in **Protocol 1000** or **2000**. Clara Beltrones in Mexico has used **Protocol 3000** quite extensively and she helped to develop it. Along with Protocol 2000, she has used it with many cancers.

ALLERGY WARNING: A very few people are allergic to **DMSO** or have very weak livers and can experience significant problems with **DMSO**. To test whether or not you are allergic to it, perform this test:

- Carefully wash and dry your arm;
- Add one drop of **DMSO** to one spot on your arm and rub it in;
- Give it about 15 minutes to soak in and then wait several hours.

If there is no pain in your liver area, it is probably safe for you to use **DMSO**, which will be the case for 99 out of 100 people. However, to be safe, wait 24 hours to make sure you have no reaction to the **DMSO**.

If you **DO** have a bad reaction to **DMSO**, you need to work on improving your liver. Begin taking a half drop of **AMS** each hour using Protocol 1000. Continue with the half drop until you feel it is safe to take more. This will clear up your liver and probably other problems that you have.

The **AMS/DMSO** treatment described below is an accelerated skin technique that pushes **AMS** into the blood plasma. In the case of cancer, it should be used in addition to a normal oral regimen of **AMS**.



DMSO Procedure Steps

1. Make an **AMS** solution by activating **20 drops of AMS** with **20 drops of Activator (4% HCl)**. Wait 30 seconds for HCl to activate. Add this solution to only one ounce of water and then spray or spread some of it liberally over one arm. (See step number 4 below for extra sick people).
2. At this point, it is best to spray 70% - 80% DMSO over the top of the **AMS** on the same area and then gently rub and spread it. If you pre-mix the **DMSO** with the **AMS**, which is practical sometimes, it will have a shelf life of about 6 hours only. You can put a plastic bag over your hand to rub the solution into your body, or you can just use your bare hands.
 - Do not use rubber gloves or other medical gloves. You could get rubber into your body as the **DMSO** could melt the rubber slightly.
 - If you notice a burning sensation, the best technique is to place a teaspoon of water on the burning area and rub it in. Keep that up until it is no longer burning.
 - You can rub olive oil and aloe vera juice on the skin after the **DMSO** application. Use a different part of the body each time you apply the **AMS/DMSO** combination. Over 8 hours, use one arm first, then the other arm, then a leg, then the other leg, then your stomach, then back to the first arm.
 - If the burning is severe, add distilled water to the **DMSO** up to 50% and more, and also to the **AMS**, as either one can cause the burning. Add enough distilled water so that the burning does not occur.
3. Repeat this process once every second hour the first day, and every hour the second and third days. Then quit for 4 days and do the same thing the next week. Remember, when you quit for 4 days, continue taking **AMS** by mouth.

After the second week you can go for 4 days a week if there is no problem with your skin.

4. For extra sick people who should take smaller doses, it would be best to start out using only a small area of the body. I suggest using an area about the size of your palm or the back of your hand, no larger. Use a small area like this for several days before going to a larger area such as the entire arm or leg. Use a larger area only if there is no bad reaction in the smaller area. Or instead, use the added distilled water trick mentioned above.

DMSO is a well-known carrier substance used widely by doctors since 1955 as a way to carry medications directly into the skin.

NOTE: Whenever drops of **AMS** are mentioned, they are activated drops. You can make activated **AMS** in using 4% HCl which is the **Activator**.